arXiv:math/0604427v1 [math.NT] 19 Apr 2006

On the Zeros of Fermat Quotients and Mirimanoff Polynomials

Björn Grohmann[1]

April 2006

Abstract

In this article an upper bound for the first consecutive zeros of the Fermat quotient is given in terms of the zeros of a Mirimanoff polynomial. This bound is obtained by investigating a relation between these polynomials and the factor system of Gauss sums of a certain number field.

Keywords: Mirimanoff polynomials, Fermat quotients, Gauss sums.

1 Introduction

Let p be an odd prime and k an integer, with $k \not\equiv 0 \bmod p$. The *Fermat quotient* $q_p(k)$ is defined to be the smallest positive integer, which satisfies the equation

$$k^{p-1} \equiv 1 + q_p(k)p \bmod p^2. \tag{1}$$

During the last century, many properties of this quotient have been investigated, mostly because of its connection to the first case of Fermat's Last Theorem: In 1909 Wieferich was able to show that if there exists an odd prime p and nonzero integers x, y, z, with $p \not| \, xyz$ and $x^p + y^p + z^p = 0$, then

$$q_p(2) = 0. \tag{2}$$

Until today, only two primes are known to satisfy equation (2): the prime 1093 (found by W. Meissner in 1913) and 3511 (N. Beeger, 1921/22).

One year after Wieferich, Mirimanoff stated that in the above case, it also holds that $q_p(3) = 0$, and it is still an open question (independent of FLT), whether there exists a prime p, such that

$$q_p(2) = q_p(3) = 0. \tag{3}$$

The interested reader should consult [4] for more on this fascinating story.

In this article we will derive a new upper bound for the integer

$$\kappa_p := \min\{n > 0 \,|\, q_p(n) \neq 0\}, \tag{4}$$

---

[1] email: nn@mhorg.de

where p denotes an odd prime. As can be easily seen from the definiton of $q_p(\cdot)$, it holds that

$$q_p(ab) \equiv q_p(a) + q_p(b) \bmod p, \tag{5}$$

so the integer κ_p is prime too.

The bound for κ_p will be in terms of the number of zeros of the *Mirimanoff polynomial*

$$\gamma_p(t) := \sum_{j=1}^{p-1} \frac{t^j}{j} \tag{6}$$

modulo p. This polynomial is closely related to the Fermat quotient, since

$$\gamma_p(t) \equiv \frac{1 - t^p - (1-t)^p}{p} \equiv (t-1)q_p(t-1) - tq_p(t) \bmod p, \tag{7}$$

and therefore

$$\kappa_p = \min\{n > 0 \,|\, \gamma_p(n) \not\equiv 0 \bmod p\}. \tag{8}$$

In the sequel we will denote the number of zeros of γ_p modulo p by $\eta_{0,p}$ and in general, for $0 \leqslant a < p$:

$$\eta_{a,p} := |\{c \mid 0 \leqslant c < p, \gamma_p(c) \equiv a \bmod p\}|. \tag{9}$$

Here, we will prove the following bound:

Theorem 1

$$\kappa_p \in O(\sqrt{\eta_{0,p}}). \tag{10}$$

We will start by recalling some basic facts about Mirimanoff polynomials and deriving some trivial bounds for κ_p. After that, we will state a connection between the factor systems of Gauss sums of a certain number field with these polynomials. This will lead to some useful relations that will enable us to prove the stated result.

2 Basic Properties

In the following, when we talk about the *zeros* of polynomial γ_p, we will always mean the zeros modulo p.

The first thing to note is that 0 and 1 are zeros of γ_p and we will call them the trivial zeros.

All the other zeros, if they exist, have multiplicity two, since $\gamma_p'(t) = 1 + t + \ldots + t^{p-2}$. From this and (8) it is immediate that

$$\kappa_p \leqslant \frac{p+1}{2}. \tag{11}$$

Looking at equation (7), we see that the Mirimanoff polynomial satisfies

$$\gamma_p(a) \equiv \gamma_p(1-a) \bmod p, \tag{12}$$

and, for $a \not\equiv 0 \bmod p$,

$$\gamma_p(a) \equiv -a\gamma_p(\frac{1}{a}) \bmod p. \tag{13}$$

From this it follows that a nontrivial zero z of γ_p always leads to the zeros:



(14)

There are two exceptional cases. In case of $\gamma_p(2) \equiv 0 \bmod p$ we have



(15)

and in case of $p \equiv 1 \bmod 3$ it is easy to see that the zeros α_6, α_6^5 of the polynomial $t^2 - t + 1 \in \mathbb{F}_p[t]$ are aswell zeros of γ_p, since



(16)

By what has been said so far, it is clear that only half of the zeros of γ_p contribute to an upper bound of κ_p, which means that

$$\kappa_p \leqslant \left\lfloor \frac{p+5}{4} \right\rfloor. \tag{17}$$

3 Relations

The Mirimanoff polynomials γ_p satisfy a number of remarkable relations. To give only one example: If z is a nontrivial zero and if in addition

$$\gamma_p(z) \equiv \gamma_p(z+1) \equiv 0 \bmod p, \tag{18}$$

then it is also true that

$$\gamma_p(z^2) \equiv 0 \bmod p. \tag{19}$$

The aim of this section is to prove the following relation.

Proposition 1 *For integers* a, b*, with* $a, b, ab \not\equiv 0, 1 \bmod p$ *we have*

$$(1-ab)\gamma_p(\frac{1-b}{1-ab}) + (1-b)\gamma_p(a) \equiv (1-ab)\gamma_p(\frac{1-a}{1-ab}) + (1-a)\gamma_p(b) \bmod p. \tag{20}$$

Here, we obtain the above example by setting $a := z$ and $b := 1/(z+1)$.

To prove the proposition, we introduce the abelian number field K, which is the real extension of the field of rationals $\mathbb{Q}$ of degree $(K : \mathbb{Q}) = p$ whose discriminant is a power of p. It may be described as the subfield $\mathbb{Q}(\Gamma)$ of the cyclotomic extension $\mathbb{Q}(\zeta_{p^2})/\mathbb{Q}$, where $\zeta_{p^2} := e^{2\pi i/p^2}$ is a primitive p^2-th root of unity, and

$$\Gamma := \mathrm{Tr}_{\mathbb{Q}(\zeta_{p^2})/K}(\zeta_{p^2}). \tag{21}$$

Hence, all nontrivial characters χ of K have conductor $f_\chi = p^2$ and are even ($\chi(-1) = 1$). As usual, we will denote the Gauss sum of a character χ of K by $\tau(\chi)$ and the elements of the factor system of these sums by

$$\omega(\chi, \chi') := \frac{\tau(\chi)\tau(\chi')}{\tau(\chi\chi')}, \tag{22}$$

for characters χ, χ' of K. For an introduction to characters of abelian number fields and Gauss sums we refer the reader to [5],[1].

We now fix a generator χ of the (cyclic) character group of K and assume that χ is normalized

$$\chi(1-p) = \zeta_p. \tag{23}$$

There is the following connection between the elements of the factor system and the Mirimonoff polynomial γ_p.

Proposition 2 *For* $k, l, k+l \not\equiv 0 \bmod p$ *it holds that*

$$\omega(\chi^k, \chi^l) = p\zeta_p^{-(k+l)\gamma_p(\frac{k}{k+l})}. \tag{24}$$

Proof. We start by evaluating the Gauss sum $\tau(\chi)$. Since $f_\chi = p^2$ and $\chi(1-p) = \zeta_p$, Theorem 1.6.2 from [1] gives

$$\tau(\chi) = p\zeta_{p^2}. \tag{25}$$

Now, for an integer k, with $(k, p) = 1$, denote by σ_k the automorphism of the Galois group $G(\mathbb{Q}(\zeta_{p^2})/\mathbb{Q})$ definied by $\zeta_{p^2}^{\sigma_k} := \zeta_{p^2}^k$. It is then well known that (cf. [2]) $\tau(\chi)^{\sigma_k} = \bar{\chi}^k(k)\tau(\chi^k)$, which leads to

$$\tau(\chi^k) = \chi^k(k)p\zeta_{p^2}^k. \tag{26}$$

The elements of the factor system may therefore be written as

$$\omega(\chi^k, \chi^l) = p\frac{\chi^k(k)\chi^l(l)}{\chi^{k+l}(k+l)}. \tag{27}$$

Since the character is normalized it follows that

$$\chi(k) = \zeta_p^{q_p(k)}, \tag{28}$$

so by (5) and (7) we obtain the stated result. □

Proof of Proposition 1. As is easily seen, the elements of the factor system satisfy the 2-cocycle condition

$$\omega(\chi, \chi'\chi'')\omega(\chi', \chi'') = \omega(\chi\chi', \chi'')\omega(\chi, \chi'), \tag{29}$$

for all characters χ, χ', χ'' of K. Using Proposition 2, this leads for integers k, l, j, with $k, l, j, k+l, l+j, k+l+j \not\equiv 0 \bmod p$, to

$$(k+l+j)\gamma_p(\frac{k}{k+l+j}) + (l+j)\gamma_p(\frac{l}{l+j}) \equiv (k+l+j)\gamma_p(\frac{k+l}{k+l+j}) + (k+l)\gamma_p(\frac{k}{k+l}).$$

We now obtain the relation of Proposition 1 by substituting $1 - a := l/(l+j)$ and $b := k/(k+l)$. □

4 Proof of the Main Theorem

As a consequence of Proposition 1 we get the following statement.

Proposition 3 *For* $1 \leqslant u, v < \kappa_p$ *it holds that*

$$\gamma_p(\frac{u}{v}) \equiv 0 \bmod p. \tag{30}$$

Proof. Deviding equation (20) by $(1-a)(1-b)$ and substituting $e := 1/(1-a)$ and $d := 1/(1-b)$ leads, for $e, d \not\equiv 0, 1 \bmod p$ and $e + d \not\equiv 1 \bmod p$, to

$$(e+d-1)\left(\gamma_p(\frac{d-1}{e+d-1}) - \gamma_p(\frac{d}{e+d-1})\right) \equiv \gamma_p(e) - \gamma_p(d) \bmod p. \tag{31}$$

The statement now easily follows by induction, taking into account the relations (12), (13) and (17). ☐

Proof of Theorem 1. To prove the bound of Theorem 1 we first note that, for any positive integer q, the integer

$$s_q := |\{(u, v) \mid 1 \leqslant u, v \leqslant q, \gcd(u, v) = 1\}| \tag{32}$$

satisfies

$$s_q \geqslant \sum_{k=1}^{q} q - q\left(\sum_{l \mid k} \frac{1}{l}\right) \geqslant q^2\left(1 - \sum_{l \leqslant q} \frac{1}{l^2}\right) \geqslant q^2\left(2 - \frac{\pi^2}{6}\right). \tag{33}$$

We now claim that for large p

$$\kappa_p \leqslant \lfloor\sqrt{p}\rfloor. \tag{34}$$

It then follows from Propostition 3 and (33) that there exist $\Omega(\kappa_p^2)$ zeros of γ_p and therefore the bound of the Theorem holds.

To prove the claim, we assume that $\kappa_p > \lfloor\sqrt{p}\rfloor$. This leads to more than

$$(\lfloor\sqrt{p}\rfloor)^2\left(2 - \frac{\pi^2}{6}\right) > \frac{p}{3} \tag{35}$$

zeros of the form a/b, with $(a, b) = 1$ and $1 \leqslant a, b \leqslant \lfloor\sqrt{p}\rfloor$. Now, if a/b is a zero of γ_p, so is $1 - a/b$. We will need the following Lemma.

Lemma 1 *Let* p *be an odd prime and* a, b *integers, with* $1 \leqslant a, b \leqslant \lfloor\sqrt{p}\rfloor$ *and* $a > b$. *Then there do* *not* *exist integers* c, d*, with* $1 \leqslant c, d \leqslant \lfloor\sqrt{p}\rfloor$*, such that:*

$$1 - \frac{a}{b} \equiv \frac{c}{d} \bmod p. \tag{36}$$

It follows that for large p and under the assumption $\kappa_p > \lfloor\sqrt{p}\rfloor$, we have produced

$$\left\lceil \frac{p}{3} \right\rceil + \left\lceil \frac{1}{2}\left(\left\lceil \frac{p}{3} \right\rceil - 1\right) \right\rceil = \frac{p+1}{2} \tag{37}$$

zeros of γ_p, which is impossible, since we did not count 0 and all nontrival zeros have multiplicity two. This means that $\kappa_p \leqslant \lfloor\sqrt{p}\rfloor$ and therefore the Theorem follows.

Proof of the Lemma. From the equation

$$bd - ad \equiv bc \bmod p \tag{38}$$

and $a > b$ it follows that there exist an integer $k > 0$, such that

$$bd - ad + kp = bc. \tag{39}$$

Now, b devides $kp - ad$, say $kp - ad = sb$, with $s > 0$ and in particular $d + s = c$, so $c > d$ and

$$1 \leqslant s \leqslant \lfloor\sqrt{p}\rfloor. \tag{40}$$

This leads to

$$kp = sb + ad < 2p, \tag{41}$$

so $k = 1$ and therefore $bd - ad + p = bc$, resp. $d = (p - bc)/(a - b)$. Since $c > d$ it follows that $ac > p$, which contradicts the assumptions of the Lemma. □

References

[1] B. C. Berndt, R. J. Evans, K. S. Williams, *Gauss and Jacobi Sums*, Can. Math. Soc. Ser. Mono. Adv. Texts, Vol 21, Wiley-Inters. Publ., 1998

[2] B. Grohmann, *Zur Arithmetik in abelschen Zahlkörpern*, Dissertation, Universität Karlsruhe, 2005, avail. at http://www.ubka.uni-karlsruhe.de/cgi-bin/psview?document=2005/informatik/25

[3] H. Hasse, *Vorlesung über Zahlentheorie*, Die Grundl. der Math. Wissensch. in Einzeld., Band LIX, Springer Berlin Göttingen Heidelberg, 1950

[4] P. Ribenboim, *13 Lectures on Fermat's Last Theorem*, Springer New York, 1979

[5] L. C. Washington, *Introduction to Cyclotomic Fields*, GTM 83, 2nd Ed., Springer New York, 1997